

dn
(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
1 February 2001 (01.02.2001)

PCT

(10) International Publication Number
WO 01/07059 A1

(51) International Patent Classification⁷: **A61K 35/12**,
35/36, 35/37, 35/38, 38/27, 38/43, 39/395, 48/00, C07H
21/04, C07K 1/00, 14/00, 16/00, 19/00, C12N 5/00, 5/02,
15/00, 15/62, 15/63, 15/85, 15/86

(21) International Application Number: PCT/US00/20055

(22) International Filing Date: 21 July 2000 (21.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/145,488 21 July 1999 (21.07.1999) US

(71) Applicant (for all designated States except US): **UNIVERSITY OF SOUTHERN CALIFORNIA** [US/US]; Suite 313, University Park, 3716 South Hope Street, Los Angeles, CA 90007 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HALL, Frederick, L.** [US/US]; - (US). **NIMNI, Marcel** [—/US]; - (US). **GORDON, Erlinda, M.** [US/US]; - (US). **BEART, Robert, W.** [—/US]; - (US).

(74) Agent: **HARRIS, Scott, C.**; Fish & Richardson P.C., Suite 500, 4350 La Jolla Village Drive, San Diego, CA 92122 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/07059 A1

(54) Title: **MATRIX-TARGETED FUSION POLYPEPTIDES FOR TISSUE REGENERATION AND WOUND HEALING**

(57) Abstract: The present invention relates to compositions and methods for promoting tissue repair and regeneration. The invention provides a fusion polypeptide useful for targeting tissues for regeneration and methods of use therefore.

**MATRIX-TARGETED FUSION POLYPEPTIDES FOR TISSUE
REGENERATION AND WOUND HEALING**

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims priority from U.S.

- 5 Provisional Application Serial No. 60/145,488, filed July
21, 1999, the disclosure of which is incorporated herein
by reference.

TECHNICAL FIELD

- 10 The present invention relates generally to
pharmaceutical agents targeted to a tissue for promoting
tissue regeneration, and more particularly to the use of
collagen-binding domains to target a growth factor to a
desired tissue.

15

BACKGROUND

- Impaired tissue healing is a significant problem in
health care. Chronic, non-healing wounds are a major
cause of prolonged morbidity in the aged human population.
- 20 Tissue repair is particularly relevant to bedridden or
diabetic patients who develop severe, external, non-
healing skin ulcers. In addition, those patients
suffering from internal lesions, such as those associated
with disorders of the digestive tract, are particularly

susceptible to the effects of non or slow-healing tissue damage.

Pharmaceutical agents that promote tissue regeneration at the site of a lesion, such as growth factors, have been utilized to accelerate wound repair. Growth factors are molecules that function not only as growth simulators (mitogens), but also as growth inhibitors. Growth factors are also known to stimulate cell migration (e.g., mitogenic cytokines), function as chemotactic agents, inhibit cell migration or invasion of tumor cells, modulate differentiated functions of cells, be involved in apoptosis, and promote survival of cells. For example, epidermal growth factor (EGF) is a mitogen that not only effects suppression of gastric acid secretion and fetal lung development, but also effects wound healing and epidermal regeneration (Franklin et al., J. Lab. Clin. Med., 108:103, 1985). EGF has been shown to be a potent stimulator of epithelial cell proliferation in the human intestine (Alison et al., Cell Biol. Int., 18:1, 1994) and other tissues.

Previous studies have demonstrated the viability of targeting extracellular matrix molecules, such as exposed collagen, for delivery of a pharmaceutical agent to a specific tissue. For example, von Willebrand Factor

(vWF)-derived collagen-binding domains have been used to target the TGF- β family of growth factors to damaged tissue (Tuan et al., Conn. Tiss. Res., 34:1, 1996; Han et al., Protein Expr. Purif., 11:169, 1997).

5 Pharmaceutical agents that promote tissue regeneration are useful not only for treating disorders associated with impaired tissue regeneration, but also for promoting tissue regeneration associated with surgical procedures, for example. Several strategies have been
10 developed to accomplish enhanced tissue repair for the treatment of damaged tissue. Within these strategies, there is a need for the controlled, sustained, site-specific targeting of a pharmaceutical agent to a wound site for the purpose of promoting tissue regeneration.

15

SUMMARY

The present invention provides new compositions and methods to induce therapeutic repair of epithelial tissue by specifically targeting tissue in need of such repair
20 with a fusion polypeptide of the invention. The invention promotes localized wound healing by providing a cell proliferation-modulating agent fused to a collagen-binding domain. The new compositions and methods are useful for wound healing in general and for promoting repair of

colonic lesions, for the capture and expansion of crypt stem cells, and the enhancement of retroviral gene transfer in colonic mucosal cells, in particular.

In one embodiment, a fusion polypeptide comprising an
5 epithelial cell proliferation-modulating agent and a collagen-binding domain capable of binding to collagen, is provided. A nucleic acid sequence encoding the fusion polypeptide is also provided.

In another aspect, the invention provides a method of
10 producing a fusion polypeptide of the invention having a collagen-binding domain and an epithelial cell proliferation-modulating agent, by growing the host cells containing a nucleic acid encoding the fusion polypeptide under conditions that allow expression of the nucleic acid
15 sequence and recovering the fusion polypeptide.

In another embodiment, the invention provides method for modulating epithelial cell proliferation in a subject. In one aspect, the modulating is by administering to the subject a fusion polypeptide comprising a collagen-binding
20 domain source linked to an epithelial cell proliferation-modulating agent. In another aspect, the modulating is by administering to the subject a therapeutically effective amount of a nucleic acid sequence encoding a fusion

polypeptide comprising a collagen-binding domain linked to an epithelial cell proliferation-modulating agent.

In a further embodiment, a tissue graft, comprising isolated tissue comprising epithelial cells treated with a fusion polypeptide comprising a collagen-binding domain linked to an epithelial cell proliferation-modulating agent, is provided. Also provided is a method of preparing a tissue graft using an effective amount of a fusion polypeptide of the invention.

10 The invention further provides a method for modulating tissue regeneration in a subject. In one aspect, the method comprises administering to the subject a therapeutically effective amount of a fusion polypeptide of the invention. In another aspect, the method comprises
15 administering to the subject a therapeutically effective amount of a nucleic acid encoding a fusion polypeptide of the invention.

In another embodiment, the invention provides a pharmaceutical composition for promoting tissue repair.

20 In one aspect, the composition comprises a fusion polypeptide comprising a collagen-binding domain linked to an epithelial cell proliferation-modulating agent in a pharmaceutically acceptable carrier. In another aspect,

the composition comprises a nucleic acid encoding a fusion polypeptide of the invention.

The details of one or more embodiments of the invention are set forth in the accompanying drawings and
5 the description below. Other features, objects, and advantages of the invention will be apparent from the description and drawings, and from the claims.

DESCRIPTION OF DRAWINGS

10 **Figure 1** is a diagram showing the structural domains of von Willebrand Factor. The A1 loop within the mature polypeptide encompasses the GPIb, collagen and heparin binding domains that function to promote platelet adhesion, collagen binding and heparin binding. The
15 minimal collagen binding sequences of human and bovine vWF, including the flanking residues, are shown.

Figure 2 is schematic diagram showing the design of recombinant EGF-CBD fusion proteins. Targeted congeners of epidermal growth factor consisting of a 6xHis
20 purification tag, an auxiliary von Willebrand factor-derived collagen-binding domain, and the cDNA sequence encoding the mature active fragment of human EGF (EGF 53+7 flanking amino acids) are shown.

Figure 3 shows the expression, induction and purification of an EGF fusion protein displaying a collagen-binding domain. Coomassie blue-stained gels identify EGF bands at molecular weight ~6kDa for both
5 collagen-targeted (EGF-CBD) and non-targeted (EGF) epidermal growth factor.

Figure 4 shows binding of EGF and EGF-CBD fusion protein to collagen matrices. Immunoreactive EGF is depicted as dark staining material in standard ELISA
10 wells, and the collagen-binding affinities were measured using a Rainbow Spectra Elisa reader and expressed as O.D. readings at A650 (plotted on the vertical axis).

Figure 5 is a graph showing the results of a biologic assay for EGF activity. The biologic activities of the
15 collagen-targeted (rEGF-CBD; black vertical bars) and non-targeted EGF (rEGF; white vertical bars) were compared to that of commercial EGF. Proliferation of NIH3T3 cells was measured by BrdU incorporation during DNA synthesis, in response to treatment with the respective EGF preparation,
20 and plotted on the vertical axis as O.D. readings at 620 nm.

Figure 6 shows *in vivo* binding of EGF-CBD to exposed collagen in a nude mouse model of ulcerative colitis. Panels A and B: Mason trichrome stain of exposed collagen

(blue-staining material) in areas of inflamed colon induced by rectal instillation of a 5% acetic acid solution. Panels C and D: Binding of EGF-CBD to exposed collagen in areas of inflamed colon as indicated by positive immunostaining (reddish-brown-staining material) using a primary antibody directed against the Hisx6 tag. Panel E: Negative binding of EGF without the collagen-binding domain; Panel F: Negative PBS control.

Figure 7 shows comparative efficacy of a collagen-targeted vs. non-targeted EGF in a mouse model of experimental colitis. The histologic features of colitis (seen as edema, hemorrhage, erosion, ulceration) three days after treatment with (Panels A-B) PBS, (Panels C-D) EGF and (Panels E-F) EGF-CBD-treated mice are shown (100X magnification).

Like reference symbols in the various drawings indicate like elements.

DETAILED DESCRIPTION

The present invention provides a recombinant fusion polypeptide comprising EGF and an appropriate collagen-binding domain (CBD) for enhancing the effective local concentration of EGF at the site of tissue injury thereby promoting repair of damaged intestinal mucosa in animal

models and, ultimately, in humans. Thus, the present invention provides a fusion polypeptide bearing an auxiliary collagen-binding domain derived from coagulation von Willebrand factor (vWF) linked to to an epithelial
5 cell proliferation-modulating agent. The invention further provides methods for using the fusion polypeptide in stimulating cell proliferation *in vitro* and promoting healing *in vivo* in an animal model.

In one embodiment, a fusion polypeptide including an
10 epithelial cell proliferation-modulating agent and a collagen-binding domain capable of binding to collagen, is provided. A nucleic acid sequence encoding the fusion polypeptide is also provided.

A "fusion polypeptide," as used herein, is a
15 polypeptide containing portions of amino acid sequence derived from two or more different proteins, or two or more regions of the same protein that are not normally contiguous. A "collagen-binding domain" is any polypeptide, or portion thereof, that can bind collagen.
20 Several collagen-binding domains are known in the art (Cruz, M.A. et al., J. Biol. Chem., 270:10822, 1995; Hoylaerts, M.F. et al., Biochem. J., 324:185, 1997; Lankhof, H. et al., Thrombos Haemostas, 75:950, 1996). In one embodiment, the collagen-binding domain is the

collagen-binding domain of von Willebrand factor, which is involved in the recognition of exposed vascular collagen (Takagi, J. et al., Biochemistry 32:8530, 1992; Tuan, T.L. et al., Conn. Tiss. Res., 34:1, 1996; Gordon, E.M. et al., Hum. Gene Ther., 8:1385, all herein incorporated by reference). von Willebrand factor was initially identified as a hemostatic factor in studies of inherited hemophilias (Wagner, Ann., Rev. Cell. Biol., 6:217, 1990), and has been shown to perform a vital surveillance function by targeting platelet aggregates to vascular lesions (Ginsburg and Bowie, Blood, 79:2507, 1992). The decapeptide WREPSFMALS (SEQ ID NO:1) has been identified to be key in the binding of von Willebrand's factor to collagen (Takagi, J. et al., supra, 1992; Tuan, T.L. et al., supra, 1996). Assays to identify collagen-binding domains of use in the subject invention are known in the art (Takagi, J. et al., supra, 1992; Tuan, T.L. et al., supra, 1996).

As used herein, an "epithelial cell proliferation-modulating agent" is any agent that can promote or inhibit epithelial cell growth or differentiation. Preferably, an epithelial cell proliferation-modulating agent of the invention is a polypeptide. More preferably, the polypeptide is a fusion polypeptide comprising a collagen-

binding domain and a growth factor, or active fragment thereof.

The term "growth factor," as used herein, includes those molecules that function as growth simulators (mitogens) or as growth inhibitors (sometimes referred to as negative growth factors). Growth factors are also known to stimulate cell migration (e.g., mitogenic cytokines), function as chemotactic agents, inhibit cell migration or invasion of tumor cells, modulate differentiated functions of cells, be involved in apoptosis, and promote survival of cells. Such factors can be secreted as diffusible factors and can also exist in membrane-anchored forms. They can, therefore, act in an autocrine, paracrine, juxtacrine, or retrocrine manner.

A cytokine is one type of growth factor. A "cytokine" is polypeptide that acts as a humoral regulator at nano-to-picomolar concentrations and which, either under normal or pathological conditions, can modulate the functional activities of individual cells and tissues. A cytokine can mediate interactions between cells directly and/or can regulate processes taking place in the extracellular environment. Cytokines comprise interleukins, lymphokines, monokines, interferons, colony-stimulating

factors, and chemokines, in addition to a variety of other proteins.

Growth factors further include epidermal growth factors (EGFs), transforming growth factors (TGFs),
5 platelet-derived growth factors (PDGFs), fibroblast growth factors (FGFs), hepatocyte growth factors (HGFs), hemopoietic growth factors (HeGFs), tumor necrosis factor (TNF-alpha), platelet-derived endothelial cell growth factor (PD-ECGF), insulin-like growth factor (IGF),
10 interleukin-8, growth hormone, angiopoietin, vascular endothelial growth factor (VEGF), acidic and basic fibroblast growth factors (FGFs), transforming growth factor alpha (TGF- α), and CYR 61 (Babic *et al.*, Proc. Natl. Acad. Sci. USA, 95:6355, 1998; Kireeva *et al.*, Mol.
15 Cell. Biol., 16:1326, 1996). Such factors further include insulin, IGF-I, IGF-II, nerve growth factor, NGF receptor, EGF, TGF- α , EGF receptor, neu, TGF- β 1, TGF- β 2, TGF- β 3, inhibin α , inhibin β , Müllerian inhibitory substance, TNF- α/β , TNF-receptor (type 1), TNF-receptor (type 2), PDGF A-chain, PDGF B-chain, PDGF receptor α , PDGF receptor β , a-FGF, b-FGF, wnt-2, hst/ks3, hepatocyte growth factor, HGF receptor (c-met),
20 IL-1 α/β , (α -chains) IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-9, IL-11, IL-12A (p35), IL-12B (p40), Interleukin 1

(type 1), Interleukin-2 α , Interleukin-2 β , Interleukin-4, Interleukin-5 α , Interleukin-6, Interleukin-7, M-CSF (also called CSF-1), M-CSF receptor (c-fms), GM-CSF, GM-CSF receptor α , GM-CSF receptor β , G-CSF, G-CSF receptor, stem cell factor, SCF receptor (c-kit), Erythropoietin (epo), epo receptor, and Leukemia inhibitory factor. Each of these molecules has been shown to induce cell proliferation, cell growth or differentiation *in vivo*. Other similar molecules that display cell growth or differentiation modulating activity are the heparin binding growth factors (HBGFs).

Epidermal growth factor (EGF) is a mitogen that can effect, for example, suppression of gastric acid secretion, fetal lung development, wound healing, and epidermal regeneration. Given the distribution and physiological responses to EGF, it appears that EGF serves to ensure the integrity and renewal of epithelial cell populations in the body. In the human gastrointestinal tract, chronic mucosal ulceration has been shown to induce the formation of a unique EGF secreting cell lineage (putative intestinal stem cells) from the base of the intestinal crypts (Wright et al., Nature, 343:82, 1990), and this EGF-secreting gland was found only in the mucosa adjacent to the site of the ulceration. Thus, EGF is

linked conceptually to colonic stem cell activity and, subsequently, to wound healing.

As used herein, the term "nucleic acid sequence" refers to a polymer of deoxyribonucleotides or ribonucleotides in the form of a separate fragment or as a component of a larger construct. Nucleic acids expressing the products of interest can be assembled from cDNA fragments or from oligonucleotides that provide a synthetic gene which is capable of being expressed in a recombinant transcriptional unit. Polynucleotide or nucleic acid sequences of the invention include DNA, RNA, and cDNA sequences.

Nucleic acid sequences utilized in the invention can be obtained by several methods. For example, the DNA can be isolated using hybridization procedures that are well known in the art. These include, but are not limited to: (1) hybridization of probes to genomic or cDNA libraries to detect shared nucleotide sequences; (2) antibody screening of expression libraries to detect shared structural features; and (3) synthesis by the polymerase chain reaction (PCR). Sequences for specific genes can also be found in GenBank, National Institutes of Health computer database.

In another aspect, the invention provides a method of producing a fusion polypeptide of the invention having a collagen-binding domain and an epithelial cell proliferation-modulating agent, by growing the host cells
5 containing a nucleic acid encoding the fusion polypeptide under conditions that allow expression of the nucleic acid sequence, and recovering the fusion polypeptide. The nucleic acid sequence of the invention can be operably linked to a promoter for expression in a prokaryotic or
10 eukaryotic expression system. For example, a nucleic acid of the invention can be incorporated in an expression vector.

Delivery of a nucleic acid of the invention can be achieved by introducing the nucleic acid into a cell using
15 a variety of methods known to those of skill in the art. For example, the construct can be delivered into a cell using a colloidal dispersion system. Alternatively, nucleic acid construct of the invention can be incorporated (*i.e.*, cloned) into an appropriate vector.
20 For purposes of expression, the nucleic acid sequences encoding the fusion polypeptide of the invention may be inserted into a recombinant expression vector. The term "recombinant expression vector" refers to a plasmid, virus, or other vehicle known in the art that has been

manipulated by insertion or incorporation of the nucleic acid sequences encoding the fusion polypeptides of the invention. The expression vector typically contains an origin of replication, a promoter, as well as specific genes that allow phenotypic selection of the transformed cells. Vectors suitable for use in the present invention include, but are not limited to, the T7-based expression vector for expression in bacteria (Rosenberg *et al.*, *Gene*, 56:125, 1987), the pMSXND expression vector for expression in mammalian cells (Lee and Nathans, *J. Biol. Chem.*, 263:3521, 1988), baculovirus-derived vectors for expression in insect cells, cauliflower mosaic virus, CaMV, tobacco mosaic virus, TMV.

Depending on the vector utilized, any of a number of suitable transcription and translation elements, including constitutive and inducible promoters, transcription enhancer elements, transcription terminators, etc. may be used in the expression vector (see, *e.g.*, Bitter *et al.*, *Methods in Enzymology*, 153:516-544, 1987). These elements are well known to one of skill in the art.

The term "operably linked" or "operably associated" refers to functional linkage between the regulatory sequence and the nucleic acid sequence regulated by the regulatory sequence. The operably linked regulatory

sequence controls the expression of the product expressed by the nucleic acid sequence. Alternatively, the functional linkage also includes an enhancer element.

"Promoter" means the minimal nucleotide sequence
5 sufficient to direct transcription. Also included in the invention are those promoter elements that are sufficient to render promoter-dependent nucleic acid sequence expression controllable for cell-type specific, tissue specific, or inducible by external signals or agents; such
10 elements may be located in the 5' or 3' regions of the native gene, or in the introns.

"Gene expression" or "nucleic acid sequence expression" means the process by which a nucleotide sequence undergoes successful transcription and
15 translation such that detectable levels of the delivered nucleotide sequence are expressed in an amount and over a time period so that a functional biological effect is achieved.

In yeast, a number of vectors containing constitutive
20 or inducible promoters may be used. (Current Protocols in Molecular Biology, Vol. 2, Ed. Ausubel et al., Greene Publish. Assoc. & Wiley Interscience, Ch. 13, 1988; Grant et al., "Expression and Secretion Vectors for Yeast," in Methods in Enzymology, Eds. Wu & Grossman, Acad. Press,

- N.Y., Vol. 153, pp.516-544, 1987; Glover, DNA Cloning, Vol. II, IRL Press, Wash., D.C., Ch. 3, 1986; "Bitter, Heterologous Gene Expression in Yeast," Methods in Enzymology, Eds. Berger & Kimmel, Acad. Press, N.Y., Vol. 5 152, pp. 673-684, 1987; and The Molecular Biology of the Yeast *Saccharomyces*, Eds. Strathern et al., Cold Spring Harbor Press, Vols. I and II, 1982). A constitutive yeast promoter, such as ADH or LEU2, or an inducible promoter, such as GAL, may be used ("Cloning in Yeast," Ch. 3, R. 10 Rothstein In: DNA Cloning Vol.11, A Practical Approach, Ed. DM Glover, IRL Press, Wash., D.C., 1986).
- Alternatively, vectors may be used which promote integration of foreign DNA sequences into the yeast chromosome.
- 15 An expression vector of the invention can be used to transform a target cell. By "transformation" is meant a permanent genetic change induced in a cell following incorporation of new DNA (i.e., DNA exogenous to the cell). Where the cell is a mammalian cell, the permanent 20 genetic change is generally achieved by introduction of the DNA into the genome of the cell. By "transformed cell" is meant a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a DNA molecule encoding a fusion protein

consisting of a collagen-binding domain linked to an epithelial cell proliferation-modulating agent, or fragment thereof. Transformation of a host cell with recombinant DNA may be carried out by conventional techniques as are well known to those skilled in the art. Where the host is prokaryotic, such as *E. coli*, competent cells which are capable of DNA uptake can be prepared from cells harvested after exponential growth phase and subsequently treated by the CaCl_2 method by procedures well known in the art. Alternatively, MgCl_2 or RbCl can be used. Transformation can also be performed after forming a protoplast of the host cell or by electroporation.

A fusion polypeptide of the invention can be produced by expression of nucleic acid encoding the protein in prokaryotes. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage DNA, plasmid DNA, or cosmid DNA expression vectors encoding a fusion protein of the invention. The constructs can be expressed in *E. coli* in large scale for *in vitro* assays. Purification from bacteria is simplified when the sequences include tags for one-step purification by nickel-chelate chromatography. The construct can also contain a tag to simplify isolation of the fusion polypeptide. For example, a polyhistidine

tag of, e.g., six histidine residues, can be incorporated at the amino terminal end of the fluorescent protein. The polyhistidine tag allows convenient isolation of the protein in a single step by nickel-chelate chromatography.

- 5 The fusion polypeptide of the invention can also be engineered to contain a cleavage site to aid in protein recovery. Alternatively, the fusion polypeptides of the invention can be expressed directly in a desired host cell for assays *in situ*.
- 10 When the host is a eukaryote, such methods of transfection of DNA as calcium phosphate co-precipitates, conventional mechanical procedures, such as microinjection, electroporation, insertion of a plasmid encased in liposomes, or virus vectors may be used.
- 15 Eukaryotic cells can also be cotransfected with DNA sequences encoding the fusion polypeptide of the invention, and a second foreign DNA molecule encoding a selectable phenotype, such as the herpes simplex thymidine kinase gene. Another method is to use a eukaryotic viral
20 vector, such as simian virus 40 (SV40) or bovine papilloma virus, to transiently infect or transform eukaryotic cells and express the protein. (Eukaryotic Viral Vectors, Cold Spring Harbor Laboratory, Gluzman ed., 1982). Preferably,

a eukaryotic host is utilized as the host cell, as described herein.

Eukaryotic systems, and preferably mammalian expression systems, allow for proper post-translational modifications of expressed mammalian proteins to occur. Eukaryotic cells that possess the cellular machinery for proper processing of the primary transcript, glycosylation, phosphorylation, and advantageously secretion of the gene product should be used as host cells for the expression of the polypeptide of the invention. Such host cell lines may include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, Jurkat, HEK-293, and WI38.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with the cDNA encoding a fusion protein of the invention controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to

form foci that, in turn, can be cloned and expanded into cell lines. For example, following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. A number of selection systems may be used, including, but not limited to, the herpes simplex virus thymidine kinase (Wigler et al., Cell, 11:223, 1977), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, Proc. Natl. Acad. Sci. USA, 48:2026, 1962), and adenine phosphoribosyltransferase (Lowy et al., Cell, 22:817, 1980) genes can be employed in tk-, hgp^rt- or ap^rt- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for dhfr, which confers resistance to methotrexate (Wigler et al., Proc. Natl. Acad. Sci. USA, 77:3567, 1980; O'Hare et al., Proc. Natl. Acad. Sci. USA, 8:1527, 1981); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, Proc. Natl. Acad. Sci. USA, 78:2072, 1981; neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., J. Mol. Biol., 150:1, 1981); and hyg^r, which confers resistance to hygromycin genes (Santerre et al., Gene, 30:147, 1984). Recently, additional selectable genes have been described, namely

- trpB, which allows cells to utilize indole in place of tryptophan; hisD, which allows cells to utilize histinol in place of histidine (Hartman & Mulligan, Proc. Natl. Acad. Sci. USA, 85:8047, 1988); and ODC (ornithine
- 5 decarboxylase), which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue L., In: Current Communications in Molecular Biology, Cold Spring Harbor Laboratory, ed., 1987).
- 10 Techniques for the isolation and purification of either microbially or eukaryotically expressed polypeptides of the invention may be by any conventional means, such as, for example, preparative chromatographic separations and immunological separations, such as those
- 15 involving the use of monoclonal or polyclonal antibodies or antigen.

A nucleic acid encoding a fusion polypeptide of the invention can also be used for gene therapy purposes. For example, when treating disorders associated with chronic

20 impaired tissue regeneration, it may be desirable to provide such tissues with a means for endogenously expressing the fusion polypeptide of the invention.

Numerous gene therapy methods that take advantage of retroviral vectors for treating a wide variety of diseases

are known in the art (see, e.g., U.S. Pat. Nos. 4,405,712
and 4,650,764; Friedmann, Science, 244:1275-1281, 1989;
Mulligan, Science, 260:926-932, 1993; and Crystal, R.,
Science 270:404-410, 1995, each of which are incorporated
5 herein by reference in their entirety). An increasing
number of these methods are currently being applied in
human clinical trials (Morgan, 1993, BioPharm, 6(1):32-35;
see also The Development of Human Gene Therapy, Theodore
Friedmann, Ed., Cold Spring Harbor Laboratory Press, Cold
10 Spring Harbor, NY, 1999; and ISBN 0-87969-528-5, which are
incorporated herein by reference in their entirety).

Another targeted delivery system useful for
introducing a nucleic acid of the invention into a target
cell is a colloidal dispersion system. Colloidal
15 dispersion systems include macromolecule complexes,
nanocapsules, microspheres, beads, and lipid-based
systems, including oil-in-water emulsions, micelles, mixed
micelles, and liposomes. The preferred colloidal system
of this invention is a liposome. Liposomes

are artificial membrane vesicles that are useful as delivery vehicles *in vitro* and *in vivo*. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0 μm can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules. RNA, DNA, and intact virions can be encapsulated within the aqueous interior and be delivered to cells in a biologically active form (Fraley et al., Trends Biochem. Sci., 6:77, 1981). In order for a liposome to be an efficient gene transfer vehicle, the following characteristics should be present: (1) encapsulation of the nucleic acid of interest (i.e., a nucleic acid encoding a fusion polypeptide of the invention or a vector comprising the nucleic acid) at high efficiency while not compromising their biological activity; (2) preferential and substantial binding to a target cell in comparison to non-target cells; (3) delivery of the aqueous contents of the vesicle to the target cell cytoplasm at high efficiency; and (4) accurate and effective expression of genetic information (Mannino et al., Biotechniques, 6:682, 1988).

In another embodiment, the invention provides method for modulating epithelial cell proliferation in a subject. In one aspect, the modulating is by administering to the

subject a fusion polypeptide comprising a collagen-binding domain source linked to an epithelial cell proliferation-modulating agent. In another aspect, the modulating is by administering to the subject a therapeutically effective
5 amount of a nucleic acid sequence encoding a fusion polypeptide comprising a collagen-binding domain linked to an epithelial cell proliferation-modulating agent. The fusion polypeptide is valuable as a therapeutic in cases in which there is impaired healing of wounds or there is a
10 need to augment normal healing mechanisms by facilitating tissue regeneration. For example, the method of the invention can be used in aiding tissue repair or regeneration at an ulcer site in a human or other subject.

"Modulation of epithelial cell proliferation" as used
15 herein, refers to regulating or controlling epithelial cell growth. The condition can include both hypertrophic (the continual multiplication of cells resulting in an overgrowth of a cell population within a tissue) and hypotrophic (a lack or deficiency of cells within a
20 tissue) cell growth or an excessive influx or migration of cells into an area of a body. The epithelial cell populations are not necessarily transformed, tumorigenic, or malignant cells, but can include normal cells as well.

The invention further provides a method for modulating tissue regeneration in a subject. In one aspect, the method comprises administering to the subject a therapeutically effective amount of a fusion polypeptide of the invention. In another aspect, the method comprises administering to the subject a therapeutically effective amount of a nucleic acid encoding a fusion polypeptide of the invention. Diseases, disorders or ailments modulated by a fusion polypeptide of the invention include tissue repair subsequent to traumatic injuries or conditions, including arthritis, osteoporosis and other skeletal disorders, and burns. Because these problems are due to a poor growth response of the fibroblasts, stem cells, chondrocytes, osteoblasts, or fibroblasts at the site of injury, the addition of an active biologic agent that stimulates or induces growth of these cells, particularly epithelial cells, is beneficial. The term "induce" or "induction" as used herein, refers to the activation, stimulation, enhancement, initiation, and/or maintenance of the cellular mechanisms or processes necessary for the formation of any of the tissue, repair process, or development, as described herein.

The compositions and methods of the invention are useful for revitalizing scar tissue resulting from

injuries due to surgical procedures, irradiation, laceration, toxic chemicals, viral infection bacterial infection, or burns. The term "scar tissue" means fibrotic or collagenous tissue formed during the healing

5 of a wound or other morbid process. For example, a fusion polypeptide of the invention can be included in a controlled release matrix that can be positioned in proximity to damaged tissue, thereby promoting regeneration of such tissue. The term "controlled release

10 matrix" means any composition that allows the slow release of a bioactive substance that is mixed or admixed therein. The matrix can be a solid composition, a porous material, or a semi-solid gel, or liquid suspension containing bioactive substances. The term "bioactive material" means

15 any composition that will modulate tissue repair when used in accordance with the method of the present invention. The bioactive materials/matrix can be introduced by means of injection, surgery, catheters, or any other means suitable for modulating tissue repair.

20 It is further envisioned that the method of the invention can be used to aid wound repair in guided tissue regeneration (GTR) procedures. Such procedures are currently used by those skilled in the medical arts to accelerate wound healing following invasive surgical

procedures. Typically, nonresorbable or bioabsorbable membranes are used to accelerate wound healing by promoting the repopulation of the wound area with cells that form the architectural and structural matrix of the tissue. For example, the method of the invention can be used in aiding periodontal tissue regeneration in a human or lower animal by placing a composition containing a bioresorbable polymer, leachable solvent, and a fusion polypeptide, comprising a collagen-binding domain and an epithelial cell proliferation-modulating agent, at a site in need of periodontal tissue regeneration in a human or other mammal such that the composition is effective for aiding tissue regeneration by releasing a therapeutically-effective amount of the fusion polypeptide at the site.

15 In another aspect, the invention is useful for the purposes of promoting tissue growth during the process of tissue engineering. By "tissue engineering" is meant the creation, design, and fabrication of biological prosthetic devices, in combination with synthetic or natural
20 materials, for the creation, augmentation, or replacement of body tissues and organs. Thus, the method can be used to augment the design and growth of tissue inside the body to repair or replace diseased or damaged tissue. A specific, non-limiting example is the use of a method of

the invention in promoting the growth of skin graft replacements that are used as a therapy in the treatment of burns and ulcers. Thus, the invention further encompasses a tissue graft, comprising epithelial cells
5 treated with a fusion polypeptide of the invention. Also provided is a method of preparing a tissue graft using an effective amount of a fusion polypeptide of the invention.

In another aspect of tissue engineering, a fusion polypeptide of the invention can be included in cell-
10 containing or cell-free devices that induce the regeneration of functional human tissues when implanted at a site that requires regeneration. As previously discussed, biomaterial-guided tissue regeneration can be used to promote epithelial cell proliferation in, for
15 example, digestive tract tissue for treatment of gastric ulcers or the pathogenic result of Krohn's disease. Thus, a fusion polypeptide of the invention can be used to promote the growth of reconstituted tissues assembled into three-dimensional configurations at the site of a wound or
20 other tissue in need of such repair.

In another aspect of tissue engineering, a fusion polypeptide of the invention can be included in external or internal devices containing human tissues designed to replace the function of diseased internal tissues. This

approach involves isolating cells from the body, placing them on or within structural matrices, and implanting the new system inside the body or using the system outside the body. The method of the invention can be included in such
5 matrices to promote the growth of tissues contained in the matrices. For example, a fusion polypeptide of the invention can be included in a cell-lined vascular graft to promote the growth of the cells contained in the graft. It is envisioned that the method of the invention can be
10 used to augment tissue repair, regeneration, and engineering in products, such as epithelial tissue, cartilage and bone, central nervous system tissues, muscle, liver, and pancreatic islet (insulin-producing) cells.

15 In another embodiment, the invention provides a pharmaceutical composition for promoting tissue repair. In one aspect, the composition comprises a fusion polypeptide, comprising a collagen-binding domain linked to an epithelial cell proliferation-modulating agent in a
20 pharmaceutically acceptable carrier. In another aspect, the composition comprises a nucleic acid encoding a fusion polypeptide of the invention.

A pharmaceutical composition according to the invention can be prepared by placing a fusion polypeptide

of the invention, or nucleic acid sequence encoding a fusion polypeptide of the invention, into a form suitable for administration to a subject using carriers, excipients, and additives or auxiliaries.

5 Generally, the terms "treating," "treatment," and the like are used herein to mean obtaining a desired pharmacologic and/or physiologic effect. The effect may be therapeutic in terms of a partial or complete cure for a cell proliferative disorder. "Treating" as used herein,
10 covers any treatment, or prevention of tissue damage, or for ameliorating the pathogenic effect of a tissue regeneration disorder, such as tissue necrosis due to diabetes, in a mammal, particularly a human, and includes:

15 (a) preventing the disease from occurring in a subject that may be predisposed to the disease, but has not yet been diagnosed as having it;

20 (b) inhibiting the disorder, i.e., arresting the development of, for example, a tumor; or

25 (c) relieving or ameliorating the disorder or disease, i.e., cause regression of the disorder or disease.

25 Thus, the invention includes various pharmaceutical compositions useful for ameliorating symptoms attributable to a tissue regeneration disorder or, alternatively, for inducing tissue regeneration following, for example, a
30 surgical procedure. Pharmaceutical compositions of the

invention are also useful for ameliorating the pathogenic effects of ulcers, for example, resulting in tissue damage.

A pharmaceutical composition according to the invention can be prepared to include a nucleic acid, or polypeptide encoded therefrom, into a form suitable for administration to a subject using carriers, excipients, and additives or auxiliaries. Frequently used carriers or auxiliaries include magnesium carbonate, titanium dioxide, lactose, mannitol and other sugars, talc, milk protein, gelatin, starch, vitamins, cellulose and its derivatives, animal and vegetable oils, polyethylene glycols and solvents, such as sterile water, alcohols, glycerol, and polyhydric alcohols. Intravenous vehicles include fluid and nutrient replenishers. Preservatives include antimicrobial, anti-oxidants, chelating agents, and inert gases. Other pharmaceutically acceptable carriers include aqueous solutions, non-toxic excipients, including salts, preservatives, buffers and the like, as described, for instance, in Remington's Pharmaceutical Sciences, 15th ed., Easton: Mack Publishing Co., 1405-1412, 1461-1487 (1975), and The National Formulary XIV., 14th ed., Washington: American Pharmaceutical Association (1975), the contents of which are hereby incorporated by

reference. The pH and exact concentration of the various components of the pharmaceutical composition are adjusted according to routine skills in the art. See Goodman and Gilman's, The Pharmacological Basis for Therapeutics (7th ed.).

The pharmaceutical compositions according to the invention may be administered locally or systemically. By "therapeutically effective dose" is meant the quantity of a compound according to the invention necessary to prevent, to cure, or at least partially arrest the symptoms of tissue damage. Amounts effective for this use will, of course, depend on the severity of the disease and the weight and general state of the patient. Typically, dosages used *in vitro* may provide useful guidance in the amounts useful for *in situ* administration of the pharmaceutical composition, and animal models may be used to determine effective dosages for treatment of particular disorders. Various considerations are described, e.g., in Langer, Science, 249: 1527, (1990); Gilman et al. (eds.) (1990), each of which is herein incorporated by reference.

As used herein, "administering a therapeutically effective amount" is intended to include methods of giving or applying a pharmaceutical composition of the invention to a subject that allow the composition to perform its

intended therapeutic function. The therapeutically effective amounts will vary according to factors, such as the degree of infection in a subject, the age, sex, and weight of the individual. Dosage regima can be adjusted
5 to provide the optimum therapeutic response. For example, several divided doses can be administered daily or the dose can be proportionally reduced as indicated by the exigencies of the therapeutic situation.

The pharmaceutical composition can be administered in
10 a convenient manner, such as by injection (subcutaneous, intravenous, etc.), oral administration, inhalation, transdermal application, or rectal administration. Depending on the route of administration, the pharmaceutical composition can be coated with a material
15 to protect the pharmaceutical composition from the action of enzymes, acids, and other natural conditions that may inactivate the pharmaceutical composition. The pharmaceutical composition can also be administered parenterally or intraperitoneally. Dispersions can also
20 be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof, and in oils. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the composition must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size, in the case of dispersion, and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols, such as mannitol, sorbitol, or sodium chloride in the

composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, aluminum monostearate and gelatin.

- 5 Sterile injectable solutions can be prepared by incorporating the pharmaceutical composition in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally,
- 10 dispersions are prepared by incorporating the pharmaceutical composition into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above.

- The pharmaceutical composition can be orally
- 15 administered, for example, with an inert diluent or an assimilable edible carrier. The pharmaceutical composition and other ingredients can also be enclosed in a hard or soft-shell gelatin capsule, compressed into tablets, or incorporated directly into the individual's
- 20 diet. For oral therapeutic administration, the pharmaceutical composition can be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and

preparations should contain at least 1% by weight of active compound. The percentage of the compositions and preparations can, of course, be varied and can conveniently be between about 5% to about 80% of the weight of the unit.

The tablets, troches, pills, capsules, and the like can also contain the following: a binder, such as gum tragacanth, acacia, corn starch, or gelatin; excipients such as dicalcium phosphate; a disintegrating agent, such as corn starch, potato starch, alginic acid, and the like; a lubricant, such as magnesium stearate; and a sweetening agent, such as sucrose, lactose or saccharin, or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it can contain, in addition to materials of the above type, a liquid carrier. Various other materials can be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules can be coated with shellac, sugar, or both. A syrup or elixir can contain the agent, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye, and flavoring, such as cherry or orange flavor. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and

substantially non-toxic in the amounts employed. In addition, the pharmaceutical composition can be incorporated into sustained-release preparations and formulations.

5 Thus, a "pharmaceutically acceptable carrier" is intended to include solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional
10 media or agent is incompatible with the pharmaceutical composition, use thereof in the therapeutic compositions and methods of treatment is contemplated. Supplementary active compounds can also be incorporated into the
15 compositions.

It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form" as used herein, refers to physically discrete units
20 suited as unitary dosages for the individual to be treated; each unit containing a predetermined quantity of pharmaceutical composition is calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for

the novel dosage unit forms of the invention are dictated by and directly dependent on: (a) the unique characteristics of the pharmaceutical composition and the particular therapeutic effect to be achieve, and (b) the
5 limitations inherent in the art of compounding such an pharmaceutical composition for the treatment of a pathogenic infection in a subject.

The principal pharmaceutical composition is compounded for convenient and effective administration in
10 effective amounts with a suitable pharmaceutically acceptable carrier in an acceptable dosage unit. In the case of compositions containing supplementary active ingredients, the dosages are determined by reference to the usual dose and manner of administration of the said
15 ingredients.

MATERIALS AND METHODS

Molecular Engineering And Cloning Of The Expression Plasmids. Figure 1 shows diagrammatically the structural
20 domains of von Willebrand Factor (vWF), identifying the primary collagen-binding domain (CBD) within the A1 loop of the mature polypeptide. The minimal collagen binding amino acid sequences of human and bovine vWF, including the flanking residues, are shown. The mature EGF

polypeptide, consisting of 53 amino acids, is generated from a large transmembrane precursor protein by proteolytic cleavage (Figure 2). In engineering the EGF-CBD fusion proteins, human coding sequences of EGF, including two additional residues at the N-terminal end and 5 amino acids at the C-terminal end of the protein were utilized. This design not only retains the original (physiological) cleavage sites, but includes these native flanking residues in an effort to facilitate the renaturation of the recombinant protein. The extended C-terminal residues (H-A-G-H-G), in particular, are considered to be important design considerations in that they are very similar to the N-terminal sequences flanking the native vWF CBD (see Figure 1). Therefore, this design is intended to optimize both the refolding of the recombinant fusion protein and the presentation of the collagen-binding domain (CBD) in solution. Moreover, the retention of the natural proteolytic cleavage site between the growth factor and the intrinsic CBD are intended to provide a mechanism for enzymatic release (i.e., "time release") of the soluble growth factor to enhance its physiological efficacy and potential therapeutic utility.

The rEGF and rEGF-CBD fusion proteins were cloned into pET expression vectors (Novagen) suitable for high-

level expression in *E. coli* (Figure 2). A strategically modified collagen-binding decapeptide derived from a functional domain within bovine von Willebrand factor (vWF; CBD; Figure 1) which normally functions in the recognition of exposed vascular collagen was utilized. A cysteine residue within the original vWF decapeptide sequence was replaced conservatively by a methionine, in order that this auxiliary domain would not interfere with the elaborate disulfide bond formation required for the folding and/or renaturation of the recombinant growth factor. Flanking linkers were also specifically designed: (i) to be devoid of native Cys residues, (ii) to include glycine residues to increase rotational flexibility and to minimize steric hindrances, and (iii) a histidine residue was included to promote an external configuration of the collagen-binding domain within the context of the homodimeric fusion protein. Thus, the design of the EGF-CBD fusion constructs, which incorporate the collagen binding decapeptide WREPSFMALS (SEQ ID NO:1) (bovine sequence) into the EGF fusion protein, was intended for targeting the biologically active growth factor to collagen exposed by injury, inflammation, ulcers, or reparative surgical procedures.

Expression, Purification, and Renaturation of regf

Fusion Proteins. The rEGF and rEGF-CBD constructs were generated from human cDNA by RT-PCR. PCR products were initially ligated into TA cloning vectors, and the

5 sequences were confirmed by direct DNA sequencing. Upon confirmation of the correct DNA sequences, the respective inserts, including linker sequences, were released by enzymatic digestion and cloned into a pET expression vector (Novagen), transformed into competent cells (BL21
10 DE3 strain of *E. coli*), and protein expression was initiated by the addition of IPTG to the culture medium. The expressed fusion proteins were isolated from *E. coli* inclusion bodies, solubilized with 8M urea, purified to near homogeneity under denaturing conditions (8M urea)
15 using nickel chelate chromatography, and renatured by oxidative refolding under optimized redox conditions.

Assessment of the Collagen-Binding Properties of regf

Fusion Proteins. The collagen-targeted or non-targeted

20 rEGF was applied onto standard ELISA plates coated with type I collagen (vitrogen-100). The plates were washed 3x with PBS, and the bound fusion protein was detected by immuno-histochemical methods, using a primary antibody directed against the Hisx6 tag (Santa Cruz SC-804) and a

HRP-labeled secondary antibody (Pierce 3146022). The colorimetric peroxidase reaction was initiated by the addition of TMB as a substrate.

- 5 **Assessment of Biological Activity of the Recombinant EGF Bearing a Collagen-binding Domain.** The mitogenic activity of the recombinant EGF fusion proteins were determined by colorimetric immunoassay of cell proliferation (Boehringer Mannheim), based on the
- 10 measurement of BrdU incorporation during DNA synthesis, and using purified commercial EGF as standardized control. Briefly, murine NIH3T3 fibroblasts (2×10^4 cells/well) in 96-well microtiter plates in 0.1 ml DMEM supplemented with 5% fetal bovine serum (D5 medium). The cells were
- 15 cultured for 3 to 4 days until confluent, at which time dilutions of samples containing EGF were added to the cell cultures, as described by George Nascimento et al., with the exception that BrdU (10 mM) is added in place of [3H] thymidine approximately 18 hours (10-24 hours) later.
- 20 After a final incubation for 24 hours, the labeling media was removed, the cells were fixed, and the incorporation of BrdU was detected by peroxidase-conjugated anti-BrdU Fab-fragments, utilizing TMB as a substrate. Each sample was assayed in triplicate wells, utilizing commercial EGF

(Sigma Chemical Co., St. Louis, MO) as a pharmacological standard. The results were quantified by measuring the absorbance at 620 nm using a scanning multiwell spectrophotometer (Phoenix).

5

In Vivo Binding of the rEGF-CBD Fusion Protein to Colonic Mucosa in a Nude Mouse Model of Experimental Colitis. In compliance with an animal protocol approved by the USC Institution Animal Care and Use Committee, six-
10 week old athymic nude mice weighing between 20-25 g were anesthetized by inhalation with methoxyflurane to effect. On Day 1, a polyetheylene tubing (4 cm long; I.D. 0.011", O.D. 0.024", Becton Dickinson, attached to a 32 G needle with 1 ml syringe) was inserted into the rectum of each
15 mouse to a depth of 2 cm. Two PBS enemas (0.5 ml each) were given to empty the colon of fecal material, after which 0.5 ml of 5% acetic acid enemas were given. The catheters were removed and the mice were then allowed to recover under a warming lamp, and returned to their cages.
20 Twenty-four hours (Day 2) after the induction of colitis, two PBS enemas were again administered to remove fecal material. Then, 0.5 ml of either PBS control, EGF or EGF-CBD (each 10 mg/ml), was given by enema. Thirty minutes later, the animals were sacrificed, the colon harvested,

and processed for immunohistochemical staining. The bound fusion protein was detected in tissue sections of colon by immunohistochemical methods, using a primary antibody directed against the Hisx6 tag and an HRP-labeled secondary antibody as described above. The colorimetric reaction was initiated by the addition of TMB as a substrate. Bound immunoreactive EGF was detected by accumulation of reddish-brown staining material on the luminal surface of injured colonic segments.

10

Efficacy of Collagen-targeted EGF (EGF-CBD) in a Nude Mouse Model of Experimental Colitis. Induction of colitis in athymic mice was conducted, as described above. For these experiments, the mice were sacrificed 24 hours (Day 3) after treatment with PBS, EGF, or EGF-CBD. The colons were extracted, washed 3 times with PBS, and fixed in 10% formalin. Tissue sections from formalin-fixed colonic segments were stained with hematoxylin-eosin or Mason trichrome stain for collagen and examined under light microscopy to determine the extent of edema, hemorrhage, inflammation, the number of well-formed colonic villi, and presence of overt stem cell proliferation. Histologic grading of severity of colitis was conducted by an

20

observer blinded to the treatment group by morphometric analysis using an Optimas image analysis system.

RESULTS

5 The present invention provides a composition and method for tethering an appropriately engineered recombinant fusion polypeptide displaying a high affinity collagen-binding domain derived from von Willebrand factor (Figure 1) linked to an epithelial cell proliferation
10 agent to sites of exposed collagen, such as those that occur at the site of a wound or lesion. The composition improves the biologic effects of, for example, EGF in an animal model of experimental colitis by cytoprotection and by modulating the proliferation of stem cells located
15 within the colonic crypts.

As shown in Figure 3, the EGF-CBD fusion protein was produced at high levels (recovery was ~80 to 100 mg /1L bacterial culture) and purified to near homogeneity by metal chelate chromatography, as determined by SDS-PAGE
20 (see Lanes 3 and 6). The fusion polypeptide can be solubilized by 6M guanidinium HCl or 8M urea, respectively, and renatured under carefully controlled redox conditions to yield a soluble, renatured polypeptide. Under optimal conditions, the bulk (> 50%)

of the purified recombinant protein was effectively renatured and recovered in soluble form. Further studies examined the physicochemical conditions of protein renaturation, including the yield (% recovery) of renatured proteins at various protein concentrations and the stabilizing effects of additives (such as sucrose or glycerol) observed upon withdrawal (dialysis) of the denaturants. These studies determined that optimal protein refolding is achieved at protein concentrations of <0.5 mg/ml, and that 20% sucrose is beneficial in optimizing the recovery of the renatured protein.

To assess the collagen-binding affinity of the rEGF-CBD fusion protein, the collagen-targeted or non-targeted rEGF was applied to collagen-coated ELISA plates and subjected to stringent washing conditions. As shown in Figure 4, the collagen-targeted EGF (EGF-CBD) exhibited enhanced binding affinity to collagen matrices compared to non-targeted EGF or PBS, demonstrating that this gain-of-function phenotype is evident and may be used to distinct advantage in animal models of wound healing. The biologic activity of the collagen-targeted rEGF-CBD, as well as the rEGF protein, was evaluated by *in vitro* cell proliferation assays using human foreskin fibroblasts and purified commercial EGF as a standardized control. Under the

experimental conditions described above by BrdU incorporation assays, dose-dependent stimulation of cell growth was observed with the commercial EGF with maximal stimulation observed at 20 ng/ml. As shown in Figure 5, the specific biological activity of each construct (rEGF and rEGF-CBD), tested at 5 ng/ml was found to be nearly 90% as active (overall mean specific activity = > 75%) as the commercial standard, indicating that the renatured EGF fusion proteins were not only refolded into soluble growth factors, but were demonstrably biologically active.

In vivo binding studies of EGF-CBD vs EGF or PBS control to exposed collagen in a nude mouse model of experimental colitis were conducted (Figure 6). Mason trichrome staining of tissue sections from colonic segments revealed significant areas of exposed collagen (Figures 6A & 6B). The EGF-CBD fusion protein was bound to and accumulated at areas of exposed collagen within colonic erosion, as indicated by positive immunostaining (brown-staining material; Figures 6C & 6D) using a primary antibody directed against the Hisx6 tag. In contrast, immunostaining of the inflamed bowel treated with non-targeted EGF (Figure 6E) or PBS (Figure 6F) was minimal to negative, indicating that the non-targeted EGF did not

adhere to the injured mucosa, and was effectively washed away by subsequent PBS infusions.

In the animal model of experimental colitis, tissue sections of colon from PBS control-treated mice (Figure 7A & B) showed extensive ulceration, and significant bleeding, edema, ulceration and acute inflammation on Day 3. While some beneficial effects were observed with the non-targeted EGF (see Figure 7C & D), the number of well-formed crypts was not significantly increased compared to the PBS group. In contrast, colonic tissue sections from EGF-CBD-treated mice (Figure 6; Table 1) showed complete regeneration of intestinal crypts, with focal areas of inflammation and ulceration and minimal bleeding. Additionally, a number of crypts showed intense stem cell proliferation, providing evidence of intestinal regeneration as well as remarkable cytoprotection.

-51-

TABLE 1

Histologic Evaluation Of Colitis By Morphometric Analysis
(Optimas)

5

Variable	None	PBS*	EGF**	EGF-CBD
Total, mm ²	3.4 ± 0.7	4.9 ± 1.0	3.0 ± 1.2 * p = 0.02	2.9 ± 0.7 * p = 0.006 ** p = 0.473
Lumen, mm ²	0.4 ± 0.2	1.6 ± 0.8	0.4 ± 0.1 * p = 0.031	0.2 ± 0.2 * p = 0.015 ** p = 0.173
Wall, mm ²	3.0 ± 0.8	3.4 ± 0.4	2.6 ± 1.2 * p = 0.142	2.7 ± 0.8 * p = 0.097 ** p = 0.441
Bleeding, %	1.6 ± 1.9	2.3 ± 2.0	0.5 ± 0.7 * p = .066	0.07 ± 0.14 * p = 0.036 ** p = 0.112
Edema, %	8.8 ± 3.3	5.6 ± 4.3	5.3 ± 3.0 * p = .446	3.9 ± 4.1 * p = 0.28 ** p = 0.310
Ulceration, %	41.3 ± 14.3	67.0 ± 39.5	33.2 ± 32.8 * p = 0.102	3.6 ± 7.2 * p = 0.012 ** p = 0.064
Inflammation, %	8.8 ± 4.8	9.9 ± 11.0	5.8 ± 6.1 * p = .249	0.2 ± 0.4 * p = 0.059 ** p = 0.059
# of Crypts	34.8 ± 11.4	24.0 ± 22.9	52.2 ± 29.8 * p = 0.085	89.2 ± 8.1 * p = 0.001 ** p = 0.027

10 * p = compared to PBS (no. of crypts: EGF-CBD > PBS:
bleeding and ulceration: EGF-CBD < PBS

** p = compared to EGF (no. of crypts: EGF-CBD > EGF);
n = 4 each group (except for PBS n = 5)

15

The studies presented herein clearly indicate that the collagen-binding function of the fusion polypeptide of the invention can effect a major and significant improvement in the rate and extent of wound healing in
5 general. The invention is useful for targeting a pharmaceutical agent to a site of epithelial tissue damage such that repair of the damaged tissue is enhanced. The invention encompasses the treatment of disorders associated with epithelial tissue damage, such as damage
10 to lung tissue or skin tissue. Though not limited to the following example, it is believed that the present fusion polypeptide will be particularly useful in the repair of digestive tract-associated lesions.

Inflammatory bowel disease, encompassing ulcerative
15 colitis and Crohn's disease (regional enteritis), are prominent causes of chronic illness in Western Europe and North America. Recently, modulation of cytokines (by IL-10 and TGF- β) that are important in the normal homeostasis of the gut immune system have been investigated in the
20 pathogenesis and treatment of experimental colitis. Conceivably, multi-modal therapies, which include anti-inflammatory activities, cytoprotective effects, and epithelial cell re-population strategies will be more

effective in attenuating or reducing the severity of inflammatory bowel disease.

EGF has been shown to be a stimulator of epithelial cell proliferation in the human intestine. In patients
5 with Zollinger-Ellison syndrome, a beneficial effect of EGF in reducing gastric secretion has been reported, although the use of EGF was not shown to be as effective as proton-pump inhibitors and histamine H2-receptor blockers. In another clinical trial for gastric ulcers,
10 patients who received intravenously administered EGF showed an increased ulcer healing rate compared to the placebo control group (Matsuo et al., Hellenic J Gastroenterol, 5(suppl 1):217, 1992). Clinical evidence of the efficacy of intravenous EGF in an infant with
15 necrotizing enterocolitis and in children with congenital microvillus atrophy has also been reported (Guglietta et al., Eur. J. Gastroenterol. Hepatol., 7:945, 1995). However, to date, there has been no clinical trials or reports of efficacy for colonic instillation of
20 recombinant EGF.

In animal models, preclinical studies on the efficacy of EGF in trinitrobenzenesulfonic acid/ethanol (TNBS)-induced colitis in immunocompetent rats have been conducted. In these studies, systemic EGF administration

reduced mucosal damage and inflammation when EGF was given intraperitoneally before, but not after, the induction of colitis, revealing a significant cytoprotective effect. In another study, systemic but not intracolonic administration of EGF was reported to accelerate the healing of colonic ulcerations in a similar model of experimental colitis. The present invention provides a novel fusion polypeptide for the local delivery of an epithelial cell proliferation-modulating agent, such as EGF, to a wound site, such as damaged intestinal epithelium, by enhanced binding of EGF to extracellular matrix molecules, specifically, to exposed collagen. Mature EGF protein was engineered to incorporate a high-affinity collagen-binding domain derived from von Willebrand factor into the primary structure of recombinant EGF fusion proteins, to specifically target EGF to sites of acute mucosal injury. This concept is based on von Willebrand factor's demonstrable surveillance function in targeting platelets to vascular lesions. Further, vWF-derived collagen-binding domains can be used to modulate the biologic activity and/or to target the TGF- β family of growth factors for specific wound healing applications.

Recently, a collagen-binding domain derived from a prokaryotic collagenase was fused to an EGF polypeptide (Nishi et al., Proc. Natl. Acad. Sci. USA, 95:7018, 1998). The recombinant fusion protein was capable of binding to exposed collagen, but was devoid of any biologic activity associated with EGF (i.e., cell proliferation modulation). The loss of biologic activity may be due to interference of the auxiliary collagen-binding domain with EGF receptor binding or with the refolding of the fusion protein during renaturation. In contrast, the EGF-CBD fusion proteins presented herein incorporate several design features that have contributed to its functional activity. The composite EGF fusion protein of the present invention embodied several aspects of molecular engineering that may indeed be critical for determining biological activity. The amino acid sequence of the fusion protein was modified by replacing several problematic cysteine residues with methionine residues (see Figure 2). In addition, flanking linkers were added to increase flexibility and to optimize the presentation of the collagen-binding domain. The aforementioned modifications yielded a fusion polypeptide with intrinsic collagen binding and growth factor activities. The EGF-CBD fusion protein was expressed at high levels in E.coli, purified to near homogeneity by

metal chelate chromatography, and was renatured by oxidative refolding into a soluble, biologically active growth factor. In solid-state assays, the collagen-targeted EGF (EGF-CBD) exhibited enhanced binding affinity to collagen matrices compared to non-targeted EGF. Further, *in vivo* binding studies in an animal model of experimental colitis showed that the collagen-targeted EGF-CBD fusion protein, but not the non-targeted EGF, and accumulated at areas of inflamed and/or ulcerated colon. Finally, the EGF-CBD fusion protein not only exhibited mitogenic activity on NIH3T3 cells *in vitro*, but also demonstrated efficacy and upon intracolonic instillation, promoted healing of the inflamed colon *in vivo* at a faster rate than non-targeted EGF. These results provide an important proof of principle that an intrinsic collagen-binding domain can be incorporated into the primary structure of EGF to regulate and, indeed, to optimize its distribution to target tissues. An appealing concept is that the display of a collagen-binding domain on the EGF protein enabled binding of the EGF fusion protein to exposed collagen at the sites of colonic erosions, and that the tethering of EGF molecules in the vicinity of colonic stem cells stimulated mitotic activity and stem cell proliferation, resulting in both cytoprotective

effects and enhanced rate of regeneration of the intestinal villi. The present invention provides a therapeutically useful fusion polypeptide for targeting damaged tissue for enhanced repair and represents a viable
5 strategy for improved delivery of a pharmaceutical agent to damaged tissue. Clinical applications of this collagen-targeted EGF fusion protein include not only the promotion of healing of colonic lesions, for example, but also stimulation of wound healing in general, capture and
10 expansion of crypt stem cells, and enhancement of retroviral gene transfer efficiency in colonic mucosal cells.

A number of embodiments of the invention have been described. Nevertheless, it will be understood that
15 various modifications may be made without departing from the spirit and scope of the invention. Accordingly, other embodiments are within the scope of the following claims.

WHAT IS CLAIMED IS:

1. A fusion polypeptide comprising a collagen-binding
domain and an epithelial cell proliferation-
5 modulating agent.
2. The fusion polypeptide of claim 1, wherein the
epithelial cell proliferation-modulating agent
stimulates epithelial cell proliferation.
- 10 3. The fusion polypeptide of claim 1, wherein the
collagen-binding domain is a collagen-binding domain
of von Willebrand factor.
- 15 4. The fusion polypeptide of claim 3, wherein the
collagen-binding domain of von Willebrand factor
comprises the decapeptide WREPSFMALS (SEQ ID NO:1).
- 20 5. The fusion protein of claim 1, wherein the epithelial
cell proliferation-modulating agent is selected from
the group consisting of a growth factor, a cytokine,
an enzyme, an enzymatic inhibitor, and an antibody.

6. The fusion polypeptide of claim 5, wherein the growth factor is selected from the group consisting of epidermal growth factor (EGF), hepatocyte growth factor (HGF), tumor necrosis factor (TNF-alpha),
5 platelet-derived endothelial cell growth factor (PD-ECGF), platelet-derived growth factor (PDGF), insulin-like growth factor (IGF), interleukin-8, growth hormone, angiopoietin, angiopoietin-1 and vascular endothelial growth factor (VEGF).
10
7. The fusion polypeptide of claim 6, wherein the growth factor is epidermal growth factor (EGF).
8. A nucleic acid sequence encoding a fusion polypeptide
15 comprising a collagen-binding domain and an epithelial cell proliferation-modulating agent.
9. The nucleic acid sequence of claim 8, operably linked to a promoter.
- 20
10. An expression vector comprising the nucleic acid sequence of claim 8.

11. The expression vector of claim 10, wherein the expression vector is a retroviral vector.
12. A host cell comprising the nucleic acid sequence of
5 claim 8.
13. A method of producing the fusion polypeptide comprising a collagen-binding domain and an epithelial cell proliferation-modulating agent,
10 comprising growing the host cells of claim 12 under conditions that allow expression of the nucleic acid sequence and recovering the fusion polypeptide.
14. The method of claim 13, wherein the host is a
15 prokaryotic cell.
15. The method of claim 13, wherein the host is a eukaryotic cell.
- 20 16. A method for modulating epithelial cell proliferation in a subject, comprising administering to the subject a fusion polypeptide comprising a collagen-binding domain linked to an epithelial cell proliferation-modulating agent.

17. The method of claim 16, wherein the subject is human.
18. The method of claim 16, wherein the subject has a
5 disorder selected from the group consisting of an
ulcerative lesion, an inflammatory lesion, a tumor,
and arthritis.
19. The fusion polypeptide of claim 16, wherein the
10 collagen-binding domain is a collagen-binding domain
of von Willebrand factor.
20. The method of claim 19, wherein the collagen-binding
domain of von Willebrand factor comprises the
15 decapeptide WREPSFMALS (SEQ ID NO:1).
21. The method of claim 16, wherein the epithelial cell
proliferation-modulating agent stimulates epithelial
cell proliferation.
- 20
22. The method of claim 16, wherein the epithelial cell
proliferation-modulating agent is selected from the
group consisting of a growth factor, a cytokine, an
enzyme, an enzymatic inhibitor, and an antibody.

23. The method of claim 22, wherein the growth factor is selected from the group consisting of epidermal growth factor (EGF), hepatocyte growth factor (HGF), tumor necrosis factor (TNF-alpha), platelet-derived endothelial cell growth factor (PD-ECGF), platelet-derived growth factor (PDGF), insulin-like growth factor (IGF), interleukin-8, growth hormone, angiopoietin, angiopoietin-1, and vascular endothelial growth factor (VEGF).
24. The method of claim 23, wherein the growth factor is epidermal growth factor (EGF).
25. A method for modulating epithelial cell proliferation in a subject, comprising administering to the subject a therapeutically effective amount of a nucleic acid sequence encoding a fusion polypeptide comprising a collagen-binding domain linked to an epithelial cell proliferation-modulating agent.
26. The method of claim 25, wherein the subject is a human.

27. The method of claim 25, wherein the subject has a disorder selected from the group consisting of an ulcerative lesion, an inflammatory lesion, a tumor, and arthritis.
- 5
28. The fusion polypeptide of claim 25, wherein the collagen-binding domain is a collagen-binding domain of von Willebrand factor.
- 10 29. The method of claim 25, wherein the collagen-binding domain of von Willebrand factor comprises the decapeptide WREPSFMALS (SEQ ID NO:1).
30. The method of claim 25, wherein the epithelial cell proliferation-modulating agent stimulates epithelial cell proliferation.
- 15
31. The method of claim 25, wherein the epithelial cell proliferation-modulating agent is selected from the group consisting of a growth factor, a cytokine, an enzyme, an enzymatic inhibitor, and an antibody.
- 20

32. The method of claim 31, wherein said cytokine is selected from the group consisting of epidermal growth factor (EGF), hepatocyte growth factor (HGF), tumor necrosis factor (TNF-alpha), platelet-derived endothelial cell growth factor (PD-ECGF), platelet-derived growth factor (PDGF), insulin-like growth factor (IGF), interleukin-8, growth hormone, angiopoietin, angiopoietin-1 and vascular endothelial growth factor (VEGF).
- 10
33. The method of claim 32, wherein the growth factor is epithelial growth factor (EGF).
34. A tissue graft, comprising isolated tissue comprising epithelial cells treated with a fusion polypeptide comprising a collagen-binding domain linked to an epithelial cell proliferation-modulating agent.
- 15
35. The tissue graft of claim 34, wherein the tissue is epithelial tissue.
- 20
36. The tissue graft of claim 34, wherein the tissue is intestinal tissue.

37. The tissue graft of claim 34, wherein the tissue is an organ.
38. The tissue graft of claim 34, wherein the collagen-binding domain is a collagen-binding domain of von Willebrand factor.
39. The tissue graft of claim 38, wherein the collagen-binding domain of von Willebrand factor comprises the decapeptide WREPSFMALS (SEQ ID NO:1).
40. The tissue graft of claim 37, wherein epithelial cell modulating agent stimulates epithelial cell proliferation.
41. The tissue graft of claim 37, wherein the epithelial cell proliferation-modulating agent is selected from the group consisting of a growth factor, a cytokine, an enzyme, an enzymatic inhibitor, and an antibody.
42. The tissue graft of claim 41, wherein the growth factor is selected from the group consisting of epidermal growth factor (EGF), hepatocyte growth factor (HGF), tumor necrosis factor (TNF-alpha),

platelet-derived endothelial cell growth factor (PD-
ECGF), platelet-derived growth factor (PDGF),
insulin-like growth factor (IGF), interleukin-8,
growth hormone, angiopoietin, angiopoietin-1 and
5 vascular endothelial growth factor (VEGF).

43. The tissue graft of claim 42, wherein the growth
factor is epithelial growth factor (EGF).

10 44. A method of preparing a tissue graft comprising
contacting isolated tissue with an effective amount
of a fusion polypeptide comprising a collagen-binding
domain linked to an epithelial cell proliferation-
modulating agent.

15

45. The method of claim 44, wherein the contacting is
in vitro.

46. The method of claim 44, wherein the contacting is
20 *in vivo*.

47. The method of claim 44, wherein the tissue is a skin.

48. The method of claim 44, wherein said tissue is an organ.
49. The method of claim 44, wherein the collagen-binding
5 domain is a von Willebrand factor collagen-binding domain.
50. The method of claim 49, wherein the collagen-binding domain of von Willebrand factor comprises the
10 decapeptide WREPSFMALS (SEQ ID NO:1).
51. The method of claim 44, wherein epithelial cell modulating agent is capable of stimulating endothelial cell proliferation.
15
52. The tissue graft of claim 37, wherein the epithelial cell proliferation-modulating agent is selected from the group consisting of a growth factor, a cytokine, an enzyme, an enzymatic inhibitor, and an antibody.
20
53. The method of claim 52, wherein the growth factor is selected from the group consisting of epidermal growth factor (EGF), hepatocyte growth factor (HGF), tumor necrosis factor (TNF-alpha), platelet-derived
25 endothelial cell growth factor (PD-ECGF), platelet-

derived growth factor (PDGF), insulin-like growth factor (IGF), interleukin-8, growth hormone, angiopoietin, angiopoietin-1 and vascular endothelial growth factor (VEGF).

5

54. The method of claim 53, wherein the growth factor is epithelial growth factor (EGF).

10

55. A method of activating a graft comprising contacting an isolated tissue with an effective amount of a nucleic acid sequence encoding a fusion polypeptide comprising a collagen-binding domain linked to an epithelial cell proliferation-modulating agent such that said nucleic acid sequence is expressed in the tissue thereby activating the graft.

15

56. The method of claim 55, wherein the nucleic acid sequence is operably linked to a promoter.

20

57. The method of claim 55, wherein the nucleic acid sequence is in an expression vector.

58. The method of claim 55, wherein the contacting is *in vitro*.

59. The method of claim 55, wherein the contacting is
in vivo.
- 5 60. The method of claim 55, wherein the tissue is
epithelial tissue.
61. The method of claim 55, wherein the tissue is the
lining of the digestive tract.
- 10 62. The method of claim 55, wherein the collagen-binding
domain is a collagen-binding domain of von Willebrand
factor.
- 15 63. The method of claim 62, wherein the collagen-binding
domain of von Willebrand factor comprises the
decapeptide WREPSFMALS (SEQ ID NO:1).
- 20 64. A pharmaceutical composition comprising a fusion
polypeptide comprising a collagen-binding domain
linked to an epithelial cell proliferation-modulating
agent in a pharmaceutically acceptable carrier.

65. A pharmaceutical composition comprising a nucleic
acid sequence encoding a fusion polypeptide
comprising a collagen-binding domain linked to
epithelial cell modulating agent in a
5 pharmaceutically acceptable carrier.





Figure 2

Molecular Engineering of Recombinant Collagen - Targeted EGF Fusion Proteins



P1 (EGF Sense Primer)

5' t ata cat atg aga aat agt gac tct gaa
M R N S D S E

P2 (EGF Antisense Primer)

5' cac gct ggc cac ggg aat tct a
3' gtg cga cag gtg ccc tta agc t
H A G H G

P3 (CBD Sense Primer)

5' tgg gag aat tct ggc cat atg tgg cgc
M W R

P4 (CBD Antisense Primer)

5' t ctg agc ggt gct ccc agt ctt ggc
3' a gac tct cca cga ggg ttc gaa cgc
L S G A P

Figur 3




Figure 4

Binding of EGF-vWF Fusion Proteins to Collagen Matrices




Figure 5






Figure 6





INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/20055

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
SWISS-PROT PROTEIN DATABASE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	HAN.B. et al. Refolding of a Recombinant Collagen-Targeted TGF-B2 Fusion Protein Expressed in Escherichia Coli. 1997. Vol. 11. pages 169-178, see entire document.	1-5, 8-10, 12-15, 65 --- 11, 17-20, 25-63
X --- Y	NISHI. N. et al. Collagen-Binding Growth Factors: Production and Characterization of Functional Fusion Proteins Having a Collagen-Binding Domain. Proc. Natl. Acad. Sci. USA. June 1998. Vol. 95. pages 7018-7023, see entire document.	1, 2, 5-10, 12-16, 21-24, 64 --- 11, 17-20, 25-63

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
I document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search
27 SEPTEMBER 2000

Date of mailing of the international search report
17 OCT 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231
Facsimile No. (703) 305-3230

Authorized official
JANET M. KERR
Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/20055

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ---- Y	GORDON. E.M. et al. Capture and Expansion of Bone Marrow-Derived Mesenchymal Progenitor Cells with a Transforming Growth Factor-B1-Von Willebrand's Factor Fusion Protein for Retrovirus-Mediated Delivery of Coagulation Factor IX. Human Gene Therapy. July 1997. Vol. 8. pages 1385-1394, see entire article.	1-5, 7-10, 12-14 ---- 11, 17-20, 25-63
Y	HALL. F.L. Targeting Retroviral Vectors to Vascular Lesions by Genetic Engineering of the MoMLV gp70 Envelope Protein. Human Gene Therapy. December 1997. Vol. 8. pages 2183-2192, see entire article.	11, 17-20, 25-63
Y	WO 98/44938 A1 (UNIVERSITY OF SOUTHER CALIFORNIA) 15 October 1998, see entire document.	11, 17-20, 25-63
Y	WO 96/25179 A1 (PURDUE RESEARCH FOUNDATION) 22 August 1996, see entire document.	35-63

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/20055

A. CLASSIFICATION OF SUBJECT MATTER: IPC (7):

A61K 35/12, 35/36, 35/37, 35/38, 38/27, 38/43, 39/395, 48/00; C07H 21/04; C07K 1/00, 14/00, 16/00, 19/00; C12N 5/00, 5/02, 15/00, 15/62, 15/63, 15/85, 15/86

A. CLASSIFICATION OF SUBJECT MATTER: US CL :

424/93.2, 93.21, 94.1, 133.1, 520, 551, 574; 435/1.1, 69.1, 70.1, 325, 440, 455, 465; 530/350, 351, 387.1, 387.3; 536/23.1, 23.2, 23.4, 23.5, 23.51, 23.53, 24.1

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

424/93.2, 93.21, 94.1, 133.1, 520, 551, 574; 435/1.1, 69.1, 70.1, 325, 440, 455, 465; 530/350, 351, 387.1, 387.3; 536/23.1, 23.2, 23.4, 23.5, 23.51, 23.53, 24.1

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

WEST, MEDLINE, EMBASE, BIOSIS, INPADOC, CAPLUS

search terms: fusion polypeptide, collagen binding domain, collagen binding domain and von Willebrand factor, cytokine, enzyme, enzymatic inhibitor, antibody, growth factor, epidermal growth factor, hepatocyte growth factor, tumor necrosis factor, platelet derived endothelial, platelet derived growth factor, insulin-like growth factor, growth hormone, angiopoietin, vascular endothelial growth factor, nucleic acid sequence and fusion polypeptide, expression vector, retroviral vector, host cell, fusion polypeptide and administration, ulcerative lesion, inflammatory lesion, tumor, arthritis, epithelial cell proliferation, tissue graft and fusion polypeptide and epithelial tissue or intestinal tissue or organ, tissue graft and nucleic acid

